

**Deposition and Survival of Enteric
Microbes in Aquatic Sediments
— A Brief Review**

Prepared with support from
Envirolink

by

Louise Weaver
Lester Sinton

Institute of Environmental Science and Research Ltd.
Christchurch

July 2009

Client Report HBRC71

**Deposition and Survival of Enteric
Microbes in Aquatic Sediments
— A Brief Review**

Dr Viv Smith

Science Programme Manager

Dr Louise Weaver

Project Leader

Dr Lester Sinton

Co-author

Dr Brent Gilpin

Peer Reviewer

DISCLAIMER

This report or document ("the Report") is given by the Institute of Environmental Science and Research Limited ("ESR") solely for the benefit of the Hawke's Bay Regional Council and other Third Party Beneficiaries as defined in the Contract between ESR and Envirolink, and is strictly subject to the conditions laid out in that Contract.

Neither ESR nor any of its employees makes any warranty, express or implied, or assumes any legal liability or responsibility for use of the Report or its contents by any other person or organisation.

ACKNOWLEDGMENTS

This report was prepared from information provided by Anna Madarasz-Smith, Hawke's Bay Regional Council. The authors gratefully acknowledge funding from the Envirolink initiative.

Critical feedback was provided by Anna Madarasz-Smith.

The authors appreciate the critical review by Dr Brent Gilpin, ESR, Christchurch.

SUMMARY	vii
1. INTRODUCTION	1
2. SETTLEMENT PROCESSES	2
3. ATTACHMENT MECHANISMS	3
4. SURVIVAL IN SEDIMENT	4
5. RELATIONSHIPS BETWEEN INDICATOR AND PATHOGEN NUMBERS IN SEDIMENTS	7
6. STRATIFICATION IN SEDIMENT	9
7. RESUSPENSION	9
8. CONCLUSIONS	11
REFERENCES	12

Summary

Hawke's Bay Regional Council (HBRC) faces problems in determining the significance of sediments as contributors to loadings of enteric indicators and pathogens in local waterways. In a study conducted by HBRC, elevated levels of faecal indicator organisms were found in recreational waters during dry weather. The study raised questions as to the origins of these elevated levels, in particular, whether they came from sediment resuspension or other sources. Accordingly, the regional council obtained an Envirolink grant to enable ESR to review this topic by:

1. Providing information on the survival of pathogens and viruses in sediments, and;
2. determining whether any information exists on the correlation between indicator bacteria and pathogens counts in sediments.

Although a review of the international literature showed that occurrence and survival of indicators and pathogens in the water column has been extensively investigated, the situation is less clear in sediments. However, there is reasonable evidence that sediments can accumulate and retain enteric indicators and pathogens. Furthermore, most published evidence suggests that survival of enteric indicators and pathogens is considerably longer in sediments than it is in the overlying water column. There is also evidence that resuspension of sediments increases microbial numbers in overlying water, particularly during storm events.

The data on the relationships between indicators and pathogens in sediments are sparse and inconclusive. Overall, there is no strong evidence of correlations between indicator and pathogen levels and survival in sediments, even though continuous deposition may be expected to have a greater integrating effect than would be expected in surface waters.

Although no evidence of increased counts of indicator and pathogenic microbes in surface waters due to resuspension was found in the HBRC studies, there was nevertheless evidence of substantially elevated counts in the sediments. There are various possible approaches to better ascertain the role of sediments in Hawke's Bay rivers and estuaries, which could include adding turbidity measurements to microbial water quality surveys, and intensive monitoring of selected sites during storm events.

1. Introduction

Three principal types of water-transmissible pathogens are found in New Zealand — bacteria, such as *Salmonella* and *Campylobacter*, protozoans (mainly *Giardia* and *Cryptosporidium*), and enteric viruses, such as the caliciviruses (which includes the norovirus group) and rotaviruses. These pathogens are costly to detect and enumerate, so their possible presence is monitored by using indicator organisms, usually *Escherichia coli* (*E. coli*) and enterococci.

The presence of both indicators and pathogens in natural waters (fresh and saline) in New Zealand has been documented by regional councils throughout the country and in nationwide surveys¹⁻⁸. However, enumeration of enteric microbes in the water column alone may underestimate the total load in a catchment, because the underlying sediments will also constitute a significant microbial reservoir. Although the presence of faecal microbes, including enteric viruses, has been reported in sediments in New Zealand coastal waters^{9, 10} and rivers¹¹, there is generally less local information available for sediments compared to that available for the overlying water bodies. Overseas studies have demonstrated the reciprocal effects of sedimentation on removing bacteria from the water column and accumulating them in bottom strata¹²⁻¹⁶.

Coastal lagoons have been assumed to protect coastal waters by attenuating pollutants¹⁶. However, there is now evidence that these coastal lagoons could act as a source of contaminants in coastal waters, because they are the recipients of and storage areas for many of the pollutants from the catchment.

In this report, we review the recent literature on the deposition and survival of key enteric indicators and pathogens in sediments. As noted above, due to a paucity of New Zealand-based information on this topic, the review relies heavily on overseas studies.

2. Settlement Processes

The extent to which indicator and pathogenic organisms are concentrated in sediments depends on the tendency of the organism to settle out of the water column, either by itself (free) or attached to particulate matter within the water.

Free bacteria can sink in a water column at rates that depend on the specific gravities of the cells, water turbulence, and their degree of motility^{12, 14, 17-19}. In most rivers, estuaries and coastal waters, natural turbulence means that settling velocities of free microbes are very slow^{20, 21}. This means deposition of individual cells will only account for a small proportion of their accumulation within sediments.

The protozoan pathogen, *Cryptosporidium*, as a free particle, has a slow settling velocity ($0.5 \mu\text{m s}^{-1}$) due to its low specific gravity²². In comparison, the larger protozoan, *Giardia*, has a faster settling velocity ($5.5 \mu\text{m s}^{-1}$).

Although there do not appear to be any published studies specifically addressing settling rates of free viruses, it is generally accepted that they can only fall out of the water column when attached to particles, as they are too small to settle naturally, even under quiescent conditions^{23, 24}.

Settling rates for enteric microbes attached to particles are much faster than when they are free entities, depending on the type and size of particle they are attached to²⁵. For example, *Giardia* and *Cryptosporidium* readily attach to particles and attachment can be as high as 93.1%^{14, 25-27}.

The ratio of attached to unattached microbes in waters varies markedly, depending on the particular organism, and this affects the rate of deposition in sediments. For instance, Characklis *et al.*¹³ studied the portions of bacteria, protozoa and viral indicators associated with settleable particles in stormwater. *E. coli* and enterococci showed a 20-35% attachment, increasing to 30-35% during storm events. The spore-forming bacterium, *Clostridium perfringens* (used as an indicator of past contamination and may be used as a surrogate for protozoan pathogens), showed the highest attachment rate at 50-70% during storm events. The rate of attachment of bacteriophages (viruses which infect bacteria, used as indicators of viral contamination) was more variable (20-60%), but was

generally found to be similar to faecal indicator bacteria. Comparable rates of attachment of *E. coli* and enterococci were reported by Fries *et al.*²⁸ (30–40%).

Enteric indicators and pathogens may be discharged into natural waters in effluent or farm runoff already attached to particulate matter^{16, 29, 30}. Others may be discharged as free entities but attach to naturally-occurring colloids and other particles in the waterway³¹. Storm events will have important, although complex, effects on this process because there will be an increase in waterway turbulence and a decrease in sedimentation. This may increase both the number of particles bearing attached indicators and the number of particles available for attachment to free microbes. The fresh input of particles from the surrounding land, and resuspension of upstream sediments will increase the sediment load during high rainfall events (see Section 7).

3. Attachment Mechanisms

Microbes attach (adsorb) to particles by a number of complex and interrelated processes. Bacteria, viruses and most other particles tend to have a net negative charge in natural waters, and thus mutually repel each other. However, processes have been identified that can overcome this repulsion. The negative charge repulsion can be overcome by the phenomenon of a compressed electric double layer³², allowing bacteria to attach to particle surfaces. Once attached, they are held in place by Van der Waals forces and hydrophobic interactions³³. This effect is most marked in waters with high ionic contents, implying that particle association is higher in saline waters^{34, 35}. Attachment by Van der Waals forces is sometimes termed weak reversible sorption, as the linkages are easily broken by shear forces such as wave action or turbulence^{36, 37}. With this type of adsorption, bacteria are not in direct contact with the particle but are held closely to the surface by the attractant forces.

A more permanent form of adsorption can also occur, which involves physical contact between bacteria and the particulate matter. It is facilitated by cellular appendages or extracellular polymers excreted from the cell³⁷. This form of attachment is not only important in removing bacteria from the water column, but may also enhance survival rates in the sediment following deposition (see Section 4).

The processes affecting virus attachment to particles have been reviewed previously^{24, 38-41}. Factors generally regarded as enhancing viral adsorption, include low pH³⁸, high ionic strength⁴², and a high cation exchange capacity⁴³ of the particle surfaces^{44, 45}.

4. Survival in Sediment

Once in the sediment, faecal indicator organisms and pathogens may remain attached to the particulate matter or they may detach. Bacteria that remain attached may survive for prolonged periods due to the associated particle offering them protection from predation and solar radiation⁴⁶. There is also evidence that some enteric bacteria, given the right conditions, can grow in sediments⁴⁷⁻⁵⁰. Although the data on the survival of viruses in sediments are sparse, there is evidence that their survival is also extended in sediments^{1, 3-5}. There is a paucity of research on the survival of protozoa in sediments but it may be assumed they will survive for long periods as there is evidence of extended survival in terrestrial environments^{12, 51, 52}.

In sediments, enteric microbes enter a very different environment to that encountered in the water column. Most importantly, when buried in bottom layers, they are protected from sunlight, which is widely recognised as the principal mechanism inactivating enteric microbes in shallow waters^{6, 7, 53}.

Most studies have suggested that the presence of organic matter in sediments significantly increases bacterial survival⁵⁴⁻⁵⁷. The organic material may provide a protective layer^{54, 58} and a supply of readily available nutrients to microbes within the sediment^{54, 59-62}. There have been, however, few correlations reported about organic matter levels and bacterial survival, therefore other factors may contribute to the overall survival of bacteria within sediments⁵⁹.

Predation and grazing may be significant factors affecting the survival of bacteria in sediments. Roper and Marshall⁶³ suggested that bacteria in sediments are protected from unicellular and multicellular predators and grazers in the interstitial spaces by two mechanisms. First, anaerobic conditions may inhibit the activities of these antagonistic organisms. Second, attachment to particles may form a physical barrier shielding bacterial cells from predation and grazing. Equally, Bauerfeind *et al.*⁶⁴ found that

bacteria survived longer in sediments sterilised by autoclaving compared with natural sediment. The increase in survival, however, may also have been due to the effect of autoclaving on sediments, which may have converted nutrients into more accessible forms for utilisation by bacteria, particularly in the absence of competitors or predators. Davies *et al.*⁶⁵ showed a net die off in bacteria, except *Clostridium perfringens* (*C. perfringens*), when predators were present. They suggested that the reduced survival in natural sediment samples was due to predation or grazing by natural protozoa.

Whatever the combination of factors that enhance survival, it is well known that bacteria persist longer in sediments than in overlying water columns^{30, 59, 63, 64}. The extent, to which enhanced survival occurs, depends on the particular bacterium. Davies *et al.*⁶⁵ found no decrease in *C. perfringens* spores in marine and freshwater sediments over a period of 28 days. In addition, the T_{90} values (time taken for 90% inactivation) for the indicator organisms, faecal coliforms and faecal streptococci, in sediment microcosms were up to 85 days. Moore *et al.* (2003)⁶⁶ found that a strain of *Salmonella* (mr-DT-104) survived for up to 119 days in sediment, which was over twice the survival rate recorded in water (54 days). Karim *et al.*⁶⁷ reported that the die-off rate of *Salmonella typhimurium* in sediment was 0.312 log per day compared to 0.345 log per day in water samples.

In general, when enteric bacteria are released into the environment, they soon succumb to the stresses they encounter, including desiccation, low nutrient levels and sunlight. However, exceptions to this have been reported. For example, Jeng *et al.*⁶⁸ conducted a laboratory-based microcosm study on the survival of *Enterococci faecalis* in sediments from an estuary in the USA. The survival curves were characterised by a growth phase followed by a stationary phase, a logarithmic decay period, and/or finally by a tailing region. Thus, an exponential decay model could not be applied to *E. faecalis* in these estuarine sediments, which appeared to demonstrate *E. faecalis* reproduction within 10 days. In addition, the authors found no evidence that survival or growth rates were affected by the amount of organic matter present in the sediment.

A recent study on freshwater sediments in Switzerland found high levels of faecal indicator organisms: 3.1×10^5 *E. coli* and 1.2×10^5 enterococci colony forming units (CFU) per 100 gram⁶⁹. In this survival experiment no faecal indicator growth was observed, possibly due to predation by naturally-occurring benthic organisms.

Far less is known about the survival of viruses in sediments compared to bacteria, and much of the literature contains conflicting information. Viruses are generally regarded as being too small to be sought as food by other microbes. They may, however, be ingested along with the particles to which they are attached, and thus be affected by the activities of interstitial grazers. Although the factors affecting virus survival in sediments do not appear to have been systematically investigated, sediment appears to enhance virus survival. For example, LaBelle and Gerba³¹ found that virus adsorption to estuarine sediment greatly increased survival time. The T_{90} (time taken for 90% reduction in live cells) value of poliovirus varied from 1.4 days in seawater alone (protected from sunlight) to 6.0 days in estuarine sediment. Increased survival of viruses (poliovirus and rotavirus) was observed in a survey of a polluted estuary in Galveston Bay⁴ where infectious viruses were isolated after 19 days when they were associated with solids, compared to up to 9 days when the viruses were freely suspended in seawater samples.

It is not possible for enteric viruses to replicate outside a host cell, so there is no possibility of growth in sediments. Bacteriophages (viruses that infect bacteria) used as surrogates for viruses were found to be more likely to adsorb to surfaces with higher total acidity and were more stable compared to free bacteriophages⁵. Although it has sometimes been suggested that enteric bacteriophages may replicate in aquatic environments^{70,71}, the evidence for their replication in natural waters is questionable^{72,73}. At present it may be assumed that bacteriophages cannot replicate in sediments.

There is also little published information on the survival of pathogenic protozoans in sediments, mostly due to the costly and laborious methods required for their culture and enumeration^{74,75}. Karim *et al.*⁶⁷ investigated the survival of *Giardia* in the water and sediments of constructed wetlands. They found that, in contrast to bacteriophages (coliphages and phage PRD1), *Giardia* survived longer in the water than in the sediment. It is important to note, however, that these experiments were conducted in microcosms in the laboratory, so the relevance of the data to the natural environment could be questioned. Like viruses, protozoan parasites can only replicate inside the body of a host, so there is no possibility of their replication in the environment. Although information on survival in sediments is limited, both *Cryptosporidium* and *Giardia* exhibit prolonged survival in other environments and this, coupled with their low infective doses, means they still pose a potential risk to humans if ingested^{76,77}.

In summary, a complex range of interacting factors appear to be involved in the survival of indicators and pathogens (and the possible growth of bacteria) in sediments. These factors are still poorly understood and more research is needed in this area.

5. Relationships Between Indicator and Pathogen Numbers in Sediments

For practical reasons, microbial water quality monitoring relies on the use of indicator organisms, such as *E. coli* and enterococci, to predict the behaviour and survival of pathogens in waterways. Although this approach can provide useful information on pathogen behaviour, recent research has highlighted problems in relying too heavily on only one or two indicator organisms to predict the behaviour of a wide range of different types of pathogens (bacterial, viral and protozoan). For example, there is now evidence that enterococci are not always reliable indicators in marine waters and waste stabilisation pond effluent^{2, 6, 7, 72}, and that they may also grow in some aquatic environments^{47, 50}. In response to this issue, a number of studies have been published on the relative survival of indicator organisms and selected pathogens in fresh and saline waters^{3, 50, 78-82}.

The relative survival rate of indicator organisms and pathogens in sediments has been investigated previously. For instance, Burton *et al.*⁸³ compared survival of the faecal indicator organism, *E. coli*, with pathogens; *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Salmonella newport* in freshwater sediments. All of the bacteria tested had prolonged survival in sediments compared to the overlying water in the microcosm used. *P. aeruginosa* and *K. pneumoniae* survived consistently longer than *E. coli* and *S. newport*. The survival of *E. coli* was comparable with *S. newport* and both survived longer in sediments with higher clay content. The authors concluded that *E. coli* was an adequate indicator of *S. newport* in several types of freshwater sediments.

Garrido-Perrez *et al.*⁷⁸ suggest that *C. perfringens* may be a good alternative to the conventional indicator organisms (namely *E. coli* and enterococci) as an indicator of faecal contamination in water and sediments as it was present in most samples and did not show any growth characteristics. In a study of an Australian estuary (water and sediment samples), *C. perfringens* was found to be the most useful indicator of faecal pollution and was the only indicator significantly correlated with the presence of pathogenic *Giardia*

and the opportunistic bacterial genus *Aeromonas*³. This study also showed that F-RNA bacteriophage was not significantly correlated with any of the pathogens examined.

Hartz *et al.*⁵⁰ investigated the survival of the faecal indicator organisms (*E. coli* and enterococci) in beach sand and experimental microcosms. Their results showed prolonged survival of faecal indicators in sediments compared with water, and there was evidence of growth in the absence of predators. The authors postulated that this may cause problems when using these organisms as faecal indicators as the high numbers present may be resuspended in the overlying water, but it is not clear that pathogens would follow the same pattern or have the ability to grow in sediments.

A study of intertidal sediments in the UK showed the absence of the pathogenic strains of *C. jejuni*, *C. coli* and *Salmonella*⁸¹. Faecal coliforms and faecal streptococci were isolated throughout the year with no obvious seasonal trends in their numbers but thermophilic *Campylobacter* spp. (not *C. jejuni* or *C. coli*) showed a strong seasonality, with greater increases in numbers occurring mostly in the summer months. *In situ* deposition of bacteria from overlying water during tidal cover showed deposition rates of approximately 0.1% of the total population of faecal coliforms, 0.01% of the faecal streptococci and 1% of the *Campylobacter* spp. in the sediments. They concluded that sediment could act as a reservoir for bacteria, especially faecal indicator organisms. However, they reported an absence of *Salmonella* and thermophilic *Campylobacter* of non-human origin present in sediments, suggesting that sediment does not act as a reservoir for these organisms.

Finally, it should be noted that it is not surprising that indicator and pathogen counts are frequently poorly correlated in the water column. Indicators, by definition, are consistently present in water polluted by faecal material; pathogens tend to be present intermittently, depending on disease incidence in the upstream communities, or carriage rates in livestock in the catchment. Furthermore, there is such a wide range of pathogen types (bacterial, viral, protozoan), that one indicator alone could not be expected to simulate the behaviour of all pathogens. Rather, there is increasing investigation into the use of “model” organisms to study the incidence and survival of pathogens, *e.g.* phage PRD1 as a model of adenoviruses. Due to the fact that sediments tend to integrate microbial pollutants, closer relationships between “similar” indicators and pathogens may possibly occur. This issue requires a lot more research before conclusions can be drawn

about the extent of similarities between pathogens and indicator organisms in sediments and their potential for resuspension.

6. Stratification in Sediment

In a water column, enteric microbes are free to circulate with water currents and turbulence. Most, however, appear to become stratified in sediments, either due to the sequence in which the particles to which they are attached are laid down, or in response to depth-related favourable and unfavourable environmental conditions. In addition, at least in freshwater sediments, some motile enteric bacteria may be able to migrate small vertical distances.

The stratification of enteric microbes in sediments does not appear to have been extensively investigated. However, in the above study of *Campylobacter* in sediments⁸¹, it was found that the faecal indicator organisms were isolated predominantly from the surface layers of the sediments and declined in number with depth, whereas *Campylobacter* spp were restricted to the surface layer. In another study, the survival of the faecal indicator organisms, *E. coli* and enterococci, was assessed in a 90-day laboratory experiment, where regular sampling was undertaken in sediment cores at different depths⁶⁹. Both *E. coli* and enterococci survived for the length of the experiment (90 days) in sediment samples close to the surface (0-6 cm depth). *E. coli* survived at all depths (0-10 cm) for up to 30 days, whereas enterococci disappeared in the deepest samples (8-10 cm) taken after 30 days.

7. Resuspension

The accumulation and enhanced survival of pathogenic bacteria and viruses in sediments (and the possible growth of some bacteria) are only important if these microbes are brought into contact with the public via direct contact or resuspension of sediments. The large number of microbes present in sediments means that even if only a small proportion of them are resuspended it may be cause for concern, for example when they come into contact with recreational water users in high enough doses to cause infection.

Enteric microbes can re-enter the water column by two mechanisms - detachment from particles, or resuspension of the particles to which they are attached. The first mechanism is probably far less important due to the fact that rapid re-attachment is likely to occur, although there is evidence that viruses can be washed from particulate matter in saline environments³⁸.

Most of the resuspension of microbes is likely to result from sediment particle resuspension, and this is in turn, most likely to result from various forms of disturbance. It has been demonstrated that tides, boats and human activity can resuspend bacteria, and to a lesser extent viruses, in the water column^{79, 84}. Even if the particles subsequently resettle, this disturbance may act to reduce microbial survival due to exposure to solar radiation in the surface waters⁸⁵. In a study of beaches in Europe, resuspension of indicator organisms only occurred in rough water, particularly during windy weather, suggesting that only sporadic resuspension was occurring⁸¹. In a survey of sediments in Switzerland, the levels of enteric microbes were found to be high, but they were not deemed high enough to cause a decline in the water quality according to EU (European Union) bathing water directives, if they were resuspended. The authors suggested that levels of 10^6 CFU per 100 g would be needed for such exceedances to occur. The same study investigated the survival times of *E. coli* in sediments in laboratory trials. Die-off of bacteria in sediments was reported to be slow, with only 10% occurring within 85 days⁶⁵. As the same proportion of *E. coli* remained culturable during the experimental period, the authors proposed that sediments offer a stable and favourable environment, which, if resuspension occurred, could provide a long-term risk of contamination. Fries *et al.*⁴⁸ reported high persistence rates in sediments of several bacteria (*E. coli*, enterococci and *Vibrio*), and that the numbers recovered significantly increased following a large storm event (Hurricane Ophelia) in the US. This research demonstrated that during heavy storms, resuspension of microbes occurred into the overlying water. Resuspension of indicator organisms and pathogens in canal sediments after heavy rainfall has also been reported in the USA⁷⁹.

8. Conclusions

Adherence to New Zealand recreational water quality guidelines only requires that *E. coli* levels in the water column are monitored. However, this may underestimate the microbial load in the catchment, because it does not account for the microbes in the sediments.

To answer the questions asked by HBRC, namely:

1. **What information is available on the survival of pathogens and viruses in sediments?**

A review of the international literature showed that there is strong evidence to suggest that sediments accumulate and retain enteric indicators and pathogens. Furthermore, most published evidence suggests that survival rates of microbes in sediments are considerably higher than in the overlying water column. Although the mechanisms are not fully understood, an increased level of organic matter appears to be the principal reason for the enhanced survival rates from the literature available.

2. **Is there any information on the correlation between sediment resuspended indicator bacteria and pathogens/viruses?**

Research in surface waters has shown that, although faecal indicator organisms fulfil a useful function in demonstrating the possible presence of pathogens and, in some cases they broadly mimic pathogen behaviour, there is a risk of over-reliance on one or two bacterial indicators.

For sediments, the data on the relationships between indicators and pathogens are sparse and inconclusive. Overall, there is no strong evidence of correlations between indicator and pathogens levels and survival in sediments, even though continuous deposition may be expected to have a greater integrating effect than would be expected in surface waters. This is an issue which requires further research before conclusions can be made.

References

- ¹ Danovaro R, Manini E, Dell'Anno A. Higher Abundance of Bacteria than of Viruses in Deep Mediterranean Sediments. *Appl Environ Microbiol.* 2002 March 1, 2002; **68**(3): 1468-72.
- ² Sinton LW, Donnison AM. Characterisation of faecal streptococci from some New Zealand effluents and receiving waters. *New Zealand Journal of Marine and Freshwater Research.* 1994; **28** 145-58.
- ³ Ferguson CM, Coote BG, Ashbolt NJ, Stevenson IM. Relationships between indicators, pathogens and water quality in an estuarine system. *Water Research.* 1996; **30**(9): 2045-54.
- ⁴ Rao VC, Seidel KM, Goyal SM, Metcalf TG, Melnick JL. Isolation of enteroviruses from water, suspended solids, and sediments from Galveston Bay: survival of poliovirus and rotavirus adsorbed to sediments. *Appl Environ Microbiol.* 1984 August 1, 1984; **48**(2): 404-9.
- ⁵ Sakoda A, Sakai Y, Hayakawa K, Suzuki M. Adsorption of viruses in water environment onto solid surfaces. *Water Science and Technology.* 1997; **35**(7): 107-14.
- ⁶ Sinton LW, Finlay RK, Lynch PA. Sunlight Inactivation of Fecal Bacteriophages and Bacteria in Sewage-Polluted Seawater. *Appl Environ Microbiol.* 1999 August 1, 1999; **65**(8): 3605-13.
- ⁷ Sinton LW, Hall CH, Lynch PA, Davies-Colley RJ. Sunlight inactivation of fecal indicator bacteria and bacteriophages from waste stabilization pond effluent in fresh and saline waters. *Appl Environ Microbiol.* 2002 Mar; **68**(3): 1122-31.

- ⁸ Till D, McBride G, Ball A, Taylor K, Pyle E. Large-scale freshwater microbiological study: rationale, results and risks. *Journal of Water and Health*. 2008 Dec; **6**(4): 443-60.
- ⁹ Lewis GD, Austin, F. J., Loutit, M., W. Enteroviruses of human origin and faecal coliforms in river water and sediments down stream from a sewage outfall in the Taieri River, Otago. *New Zealand Journal of Marine and Freshwater Research*. 1986; **20**: 101-5.
- ¹⁰ Loutit MW, Lewis, G. D. Faecal bacteria from sewage effluent in sediments around a sewage outfall. *New Zealand Journal of Marine and Freshwater Research*. 1985; **19**: 177-85.
- ¹¹ Muirhead RW, Davies-Colley RJ, Donnison AM, Nagels JW. Faecal bacteria yields in artificial flood events: quantifying in-stream stores. *Water Res*. 2004 Mar; **38**(5): 1215-24.
- ¹² Brookes JD, Antenucci J, Hipsey M, Burch MD, Ashbolt NJ, Ferguson C. Fate and transport of pathogens in lakes and reservoirs. *Environment International*. 2004; **30**(5): 741-59.
- ¹³ Characklis GW, Dilts MJ, Simmons Iii OD, Likirdopoulos CA, Krometis L-AH, Sobsey MD. Microbial partitioning to settleable particles in stormwater. *Water Research*. 2005; **39**(9): 1773-82.
- ¹⁴ Dai X, Boll J. Settling velocity of *Cryptosporidium parvum* and *Giardia lamblia*. *Water Research*. 2006; **40**(6): 1321-5.
- ¹⁵ Deloffre J, Lafite R, Lesueur P, Lesourd S, Verney R, Guézennec L. Sedimentary processes on an intertidal mudflat in the upper macrotidal Seine estuary, France. *Estuarine, Coastal and Shelf Science*. 2005; **64**(4): 710-20.

- ¹⁶ Steets BM, Holden PA. A mechanistic model of runoff-associated fecal coliform fate and transport through a coastal lagoon. *Water Research*. 2003; **37**(3): 589-608.
- ¹⁷ Jassby AD. The ecological significance of sinking to planktonic bacteria. *Canadian Journal of Microbiology*. 1975; **21**: 270-4.
- ¹⁸ Pedrós-Alió C, Mas J, Gasol JM, Guerrero R. Sinking speeds of free-living phototrophic bacteria determined with covered and uncovered traps. *J Plankton Res*. 1989 January 1, 1989; **11**(5): 887-905.
- ¹⁹ Bailey MC, Hamilton DP. Wind induced sediment resuspension: a lake-wide model. *Ecological Modelling*. 1997; **99**(2-3): 217-28.
- ²⁰ Bai S, Lung W-S. Modeling sediment impact on the transport of fecal bacteria. *Water Research*. 2005; **39**(20): 5232-40.
- ²¹ Mahler BJ, Personné JC, Lods GF, Drogue C. Transport of free and particulate-associated bacteria in karst. *Journal of Hydrology*. 2000; **238**(3-4): 179-93.
- ²² Gray NF. Monitoring and removal of pathogens. *Drinking Water Quality: Problems and Solutions*. 2nd ed: Cambridge University Press 2008:316-7.
- ²³ Yavuz Corapcioglu M, Haridas A. Microbial transport in soils and groundwater: A numerical model. *Advances in Water Resources*. 1985; **8**(4): 188-200.
- ²⁴ Yates MV, Yates, S. R. Virus survival and transport in groundwater. *Water Science and Technology*. 1988; **20**: 301-7.
- ²⁵ Searcy KE, Packman AI, Atwill ER, Harter T. Association of *Cryptosporidium parvum* with Suspended Particles: Impact on Oocyst Sedimentation. *Appl Environ Microbiol*. 2005 February 1, 2005; **71**(2): 1072-8.

- ²⁶ Kuczynska E, Shelton DR, Pachepsky Y. Effect of Bovine Manure on *Cryptosporidium parvum* Oocyst Attachment to Soil. *Appl Environ Microbiol.* 2005 October 1, 2005; **71**(10): 6394-7.
- ²⁷ Medema GJ, Schets FM, Teunis PFM, Havelaar AH. Sedimentation of Free and Attached *Cryptosporidium* Oocysts and *Giardia* Cysts in Water. *Appl Environ Microbiol.* 1998 November 1, 1998; **64**(11): 4460-6.
- ²⁸ Fries JS, Characklis, G. W., Noble, R. T. Attachment of Fecal Indicator Bacteria to Particles in the Neuse River Estuary, N.C. *Journal of Environmental Engineering.* 2006; **132**: 1338-45.
- ²⁹ Servais P, Garcia-Armisen T, George I, Billen G. Fecal bacteria in the rivers of the Seine drainage network (France): Sources, fate and modelling. *Science of The Total Environment.* 2007; **375**(1-3): 152-67.
- ³⁰ Van Donsel DJ, Geldreich EE, Clarke NA. Seasonal Variations in Survival of Indicator Bacteria in Soil and Their Contribution to Storm-water Pollution. *Appl Environ Microbiol.* 1967 November 1, 1967; **15**(6): 1362-70.
- ³¹ LaBelle R, Gerba CP. Investigations into the protective effect of estuarine sediment on virus survival. *Water Research.* 1982; **16**(4): 469-78.
- ³² Stumm W, Morgan, J. J. *Aquatic Chemistry.* 2nd ed. New York: Wiley-Interscience 1981.
- ³³ Tsao YH, Evans DF, Wennerstroem H. Long-range attraction between a hydrophobic surface and a polar surface is stronger than that between two hydrophobic surfaces. *Langmuir.* 1993 03/01; **9**(3): 779-85.
- ³⁴ Gannon JT, Mingelgrin U, Alexander M, Wagenet RJ. Bacterial transport through homogeneous soil. *Soil Biology and Biochemistry.* 1991; **23**(12): 1155-60.

- ³⁵ Tan Y, Cannon, J.T. , Baveye, P., Alexander, M. 1994. Water Resour. Res. 30:3243–3252. . Transport of bacteria in an aquifer sand—Experiments and model simulations. *Water Resource Research*. 1994; **30**: 3243-52.
- ³⁶ Boks NP, Norde W, van der Mei HC, Busscher HJ. Forces involved in bacterial adhesion to hydrophilic and hydrophobic surfaces. *Microbiology*. 2008 October 1, 2008; **154**(10): 3122-33.
- ³⁷ Palmateer GA, McLean, D. E., Kutas, W. L., Meissner, S. M. Suspended particulate/bacterial interaction in agricultural drains. In: Rao SS, ed. *Particulate matter and aquatic contaminants*. Florida: Boca Raton 1993:1-40.
- ³⁸ Gerba CP, Stagg CH, Abadie MG. Characterization of sewage solid-associated viruses and behavior in natural waters. *Water Research*. 1978; **12**(10): 805-12.
- ³⁹ Yates MV, Gerba, C. P., Kelly, L. M. Virus persistence in groundwater. *Applied and Environmental Microbiology*. 1985; **49**: 778-81.
- ⁴⁰ Yates MV, Yates, S. R. A comparison of geostastical methods for estimating virus inactivation rates in ground water. *Water Research*. 1987; **21**: 1119-25.
- ⁴¹ Mathess G, Pekdeger, A., Schroeter, J. Persistence and transport of bacteria and viruses in groundwater — a conceptual evaluation. *Journal of Contaminant Hydrology*. 1988; **2**: 171-88.
- ⁴² Redman JA, Walker SL, Elimelech M. Bacterial Adhesion and Transport in Porous Media: Role of the Secondary Energy Minimum. *Environmental Science & Technology*. 2004; **38**(6): 1777-85.

- ⁴³ Lipson SM, Stotzky G. Adsorption of reovirus to clay minerals: effects of cation-exchange capacity, cation saturation, and surface area. *Appl Environ Microbiol.* 1983 September 1, 1983; **46**(3): 673-82.
- ⁴⁴ Moore BE, Sagik, B. P., Sorber, C. A. Viral transport to groundwater at a wastewater land application site. *Journal of Water Pollution Control Federation.* 1981; **53**: 1492-502.
- ⁴⁵ Vilker VL, Fong, J. C., Seyed-Hoseyni, M. Poliovirus adsorption to narrow particle size fractions of sand and montmorillonite clay. *Journal of Colloid Interface Science.* 1983; **92**: 422-35.
- ⁴⁶ Anderson KL, Whitlock JE, Harwood VJ. Persistence and Differential Survival of Fecal Indicator Bacteria in Subtropical Waters and Sediments. *Appl Environ Microbiol.* 2005 June 1, 2005; **71**(6): 3041-8.
- ⁴⁷ Desmarais TR, Solo-Gabriele HM, Palmer CJ. Influence of Soil on Fecal Indicator Organisms in a Tidally Influenced Subtropical Environment. *Appl Environ Microbiol.* 2002 March 1, 2002; **68**(3): 1165-72.
- ⁴⁸ Fries JS, Characklis GW, Noble RT. Sediment-water exchange of *Vibrio* sp. and fecal indicator bacteria: Implications for persistence and transport in the Neuse River Estuary, North Carolina, USA. *Water Research.* 2008; **42**(4-5): 941-50.
- ⁴⁹ Grossart H-P, Kiorboe T, Tang K, Ploug H. Bacterial Colonization of Particles: Growth and Interactions. *Appl Environ Microbiol.* 2003 June 1, 2003; **69**(6): 3500-9.
- ⁵⁰ Hartz A, Cuvelier M, Nowosielski K, Bonilla TD, Green M, Esiobu N, et al. Survival Potential of *Escherichia coli* and Enterococci in Subtropical Beach Sand: Implications for Water Quality Managers. *J Environ Qual.* 2008 May 1, 2008; **37**(3): 898-905.

- 51 Ives RL, Kamarainen AM, John DE, Rose JB. Use of Cell Culture To Assess *Cryptosporidium parvum* Survival Rates in Natural Groundwaters and Surface Waters. *Appl Environ Microbiol.* 2007 September 15, 2007; **73**(18): 5968-70.
- 52 Robertson LJ, Campbell AT, Smith HV. Survival of *Cryptosporidium parvum* oocysts under various environmental pressures. *Appl Environ Microbiol.* 1992 November 1, 1992; **58**(11): 3494-500.
- 53 Davies-Colley RJ, Bell RG, Donnison AM. Sunlight Inactivation of Enterococci and Fecal Coliforms in Sewage Effluent Diluted in Seawater. *Appl Environ Microbiol.* 1994 June 1, 1994; **60**(6): 2049-58.
- 54 Brown CM, Eilwood, D. C., Hunter, J. R. 1977. Growth of bacteria at surfaces: Influence of nutrient concentrations. *FEMS Microbiology Letters.* 1977; **1**: 163-6.
- 55 Gerba CP, McLeod, J. S. Effect of sediments on the survival of *Escherichia coli* in marine waters. *Applied and Environmental Microbiology.* 1976; **32**(1): 114-20.
- 56 Hood MA, Ness GE. Survival of *Vibrio cholerae* and *Escherichia coli* in estuarine waters and sediments. *Appl Environ Microbiol.* 1982 March 1, 1982; **43**(3): 578-84.
- 57 Ghoul M, Bernard T, Cormier M. Evidence that *Escherichia coli* accumulates glycine betaine from marine sediments. *Appl Environ Microbiol.* 1990 February 1, 1990; **56**(2): 551-4.
- 58 Gauthier MJ, Le Rudulier D. Survival in seawater of *Escherichia coli* cells grown in marine sediments containing glycine betaine. *Appl Environ Microbiol.* 1990 September 1, 1990; **56**(9): 2915-8.
- 59 Shiaris MP, Rex AC, Pettibone GW, Keay K, McManus P, Rex MA, et al. Distribution of indicator bacteria and *Vibrio parahaemolyticus* in sewage-polluted intertidal sediments. *Appl Environ Microbiol.* 1987 August 1, 1987; **53**(8): 1756-61.

- ⁶⁰ Milne DP, Curran, J. C., Findlay, J. S., Crowther, J. M., Bennett, J., Wood, B. J. B. The effect of dissolved nutrients and inorganic suspended solids on the survival of *E.coli* in seawater. *Water Science and Technology*. 1991; **24**: 133-6.
- ⁶¹ Luna GM, Manini E, Danovaro R. Large Fraction of Dead and Inactive Bacteria in Coastal Marine Sediments: Comparison of Protocols for Determination and Ecological Significance. *Appl Environ Microbiol*. 2002 July 1, 2002; **68**(7): 3509-13.
- ⁶² Rusch A, Forster S, Huettel M. Bacteria, diatoms and detritus in an intertidal sandflat subject to advective transport across the water-sediment interface. *Biogeochemistry*. 2001; **55**(1): 1-27.
- ⁶³ Roper MM, Marshall KC. Biological Control Agents of Sewage Bacteria in Marine Habitats. *Marine and Freshwater Research*. 1978; **29**(3): 335-43.
- ⁶⁴ Bauerfeind S, Gerhardt, G. G., Rheinheimer, G. Investigations on the survival of faecal bacteria in experiments with and without sediment addition. *Zbl Bakt Hyg, I Abt Orig B*. 1981; **174**: 364-74.
- ⁶⁵ Davies CM, Long JA, Donald M, Ashbolt NJ. Survival of fecal microorganisms in marine and freshwater sediments. *Appl Environ Microbiol*. 1995 May 1, 1995; **61**(5): 1888-96.
- ⁶⁶ Moore BC, Martinez E, Gay JM, Rice DH. Survival of *Salmonella enterica* in Freshwater and Sediments and Transmission by the Aquatic Midge *Chironomus tentans* (Chironomidae: Diptera). *Appl Environ Microbiol*. 2003 August 1, 2003; **69**(8): 4556-60.
- ⁶⁷ Karim MR, Manshadi FD, Karpiscak MM, Gerba CP. The persistence and removal of enteric pathogens in constructed wetlands. *Water Research*. 2004; **38**(7): 1831-7.

- 68 Jeng HAC, Sinclair, R., Daniels, R., Andrew, E. Survival of *Enterococci faecalis* in estuarine sediments. *International Journal of Environmental Studies*. 2005; **62**: 283-91.
- 69 Haller L, Poté J, Loizeau J-L, Wildi W. Distribution and survival of faecal indicator bacteria in the sediments of the Bay of Vidy, Lake Geneva, Switzerland. *Ecological Indicators*. **In Press, Corrected Proof**.
- 70 Vaughn JM, Metcalf, T. G. Coliphages as indicators of enteric viruses in shellfish and shellfish raising estuarine waters. *Water Research*. 1975; **9**(7): 613-6.
- 71 Borrego JJ, Cornax, R., Morinigo, M.A., Martinez-Manzares, E. Romero, P. Coliphages as an indicator of faecal pollution in water. *Water Research*. 1990; **24**: 111-6.
- 72 Sinton LW. Survival of enteric bacteria in seawater: Biotic and abiotic factors. In: Belkin SS, Colwell, R. R., ed. *Oceans and Health: Pathogens in the Marine Environment*. New York: Kluwer Academic/Plenum Publishers 2005.
- 73 Hernandez-Delgado EA, Toranzos, G. A. *In-situ* replication studies of somatic and male-specific coliphages in a tropical pristine river. *Water Science and Technology*. 1995; **31**(5-6): 247-50.
- 74 Bradford SA, Bettahar M. Straining, Attachment, and Detachment of *Cryptosporidium* Oocysts in Saturated Porous Media. *J Environ Qual*. 2005 March 1, 2005; **34**(2): 469-78.
- 75 Bradford SA, Tadassa YF, Pachepsky Y. Transport of *Giardia* and Manure Suspensions in Saturated Porous Media. *J Environ Qual*. 2006 April 3, 2006; **35**(3): 749-57.

- ⁷⁶ DuPont HL, Chappell CL, Sterling CR, Okhuysen PC, Rose JB, Jakubowski W. The Infectivity of *Cryptosporidium parvum* in Healthy Volunteers. *N Engl J Med*. 1995 March 30, 1995; **332**(13): 855-9.
- ⁷⁷ Rose JB, Haas CN, Regli S. Risk assessment and control of waterborne giardiasis. *Am J Public Health*. 1991 June 1, 1991; **81**(6): 709-13.
- ⁷⁸ Garrido-Pérez MC, Anfuso E, Acevedo A, Perales-Vargas-Machuca JA. Microbial indicators of faecal contamination in waters and sediments of beach bathing zones. *International Journal of Hygiene and Environmental Health*. 2008; **211**(5-6): 510-7.
- ⁷⁹ Goyal SM, Gerba CP, Melnick JL. Occurrence and distribution of bacterial indicators and pathogens in canal communities along the Texas coast. *Appl Environ Microbiol*. 1977 August 1, 1977; **34**(2): 139-49.
- ⁸⁰ Lisle JT, Smith JJ, Edwards DD, McFeters GA. Occurrence of Microbial Indicators and *Clostridium perfringens* in Wastewater, Water Column Samples, Sediments, Drinking Water, and Weddell Seal Feces Collected at McMurdo Station, Antarctica. *Appl Environ Microbiol*. 2004 December 1, 2004; **70**(12): 7269-76.
- ⁸¹ Obiri-Danso K, Jones K. Intertidal sediments as reservoirs for hippurate negative campylobacters, salmonellae and faecal indicators in three EU recognised bathing waters in North West England. *Water Research*. 2000; **34**(2): 519-27.
- ⁸² Touron A, Berthe T, Gargala G, Fournier M, Ratajczak M, Servais P, et al. Assessment of faecal contamination and the relationship between pathogens and faecal bacterial indicators in an estuarine environment (Seine, France). *Marine Pollution Bulletin*. 2007; **54**(9): 1441-50.
- ⁸³ Burton GA, Jr., Gunnison D, Lanza GR. Survival of pathogenic bacteria in various freshwater sediments. *Appl Environ Microbiol*. 1987 April 1, 1987; **53**(4): 633-8.

- ⁸⁴ Pettibone GW, Irvine KN, Monahan KM. Impact of a ship passage on bacteria levels and suspended sediment characteristics in the Buffalo River, New York. *Water Research*. 1996; **30**(10): 2517-21.
- ⁸⁵ Simental L, Martinez-Urtaza J. Climate Patterns Governing the Presence and Permanence of Salmonellae in Coastal Areas of Bahia de Todos Santos, Mexico. *Appl Environ Microbiol*. 2008 October 1, 2008; **74**(19): 5918-24.